

Name _____

Math 312 - Section 002 - Midterm 1 (Practice exam)
Thursday, February 14, 2019, @ 10:30 AM - 11:50 AM

No form of cheating will be tolerated. You are expected to uphold the Code of Academic Integrity. I certify that all of the work on this test is my own.

Signature: _____

The use of calculators, computers and similar devices is neither necessary nor permitted during this exam. Correct answers without proper justification will not receive full credit. Clearly highlight your answers and the steps taken to arrive at them: illegible work will not be graded. You may use both sides of one 8×11 cheat-sheet.

OFFICIAL USE ONLY:

Problem	Points	Your score
1	20	
2	20	
3	20	
4	20	
5	20	
Total	100	

Problem 1 [20 points]

Consider the matrix $A = \begin{bmatrix} 1 & 3 & 1 & 1 & 1 \\ 0 & 0 & -1 & -1 & 0 \\ 2 & 6 & 4 & 4 & 3 \\ 0 & 0 & 2 & 2 & 3 \end{bmatrix}$.

Part a. Find the matrix R , the RREF (row reduced echelon form) of A .

Part b. Find the matrix E that transforms A into R , i.e., $EA = R$.

Part c. Find the complete solution to $A\mathbf{x} = \begin{bmatrix} 0 \\ 0 \\ 1 \\ 3 \end{bmatrix}$.

Part d. Find a basis for each one of the four fundamental subspaces of A .

Problem 2 [20 points]

Forward elimination changes $A\vec{x} = \vec{b}$ to a row reduced echelon form $R\vec{x} = \vec{d}$. The complete solution is

$$\vec{x} = \begin{bmatrix} 4 \\ 0 \\ 0 \end{bmatrix} + \alpha \begin{bmatrix} 2 \\ 1 \\ 0 \end{bmatrix} + \beta \begin{bmatrix} 5 \\ 0 \\ 1 \end{bmatrix}, \quad \alpha, \beta \in \mathbb{R}.$$

Part a. What is the 3 by 3 reduced row echelon matrix R and what is \vec{d} ?

Part b. If the process of elimination subtracted 3 times row 1 from row 2 and then 5 times row 1 from row 3, what matrix connects R and \vec{d} to the original A and \vec{b} ?

Part c. Find the LU decomposition of A . (Hint: use part b)

Problem 3 [20 points]

$$\text{Let } \mathbf{u}_1 = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}, \mathbf{u}_2 = \begin{bmatrix} 2 \\ 1 \\ 0 \end{bmatrix}, \mathbf{u}_3 = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \mathbf{u}_4 = \begin{bmatrix} 2 \\ 0 \\ 2 \end{bmatrix}, \mathbf{u}_5 = \begin{bmatrix} 0 \\ 2 \\ 1 \end{bmatrix}.$$

Part a. Find a basis for $V = \{\vec{u}_1, \vec{u}_2, \vec{u}_3, \vec{u}_4, \vec{u}_5\}$. What is the dimension of V ?

Part b. Show that $\{1, 1 - x, (1 - x)^2\}$ is a basis of \mathcal{P}_2 , the vector space of polynomials of degree at most 2.

Problem 4 [20 points]

Part a. Given the following bases for the space of polynomials of degree at most 2, \mathcal{P}_2 , find $P_{\mathcal{C} \leftarrow \mathcal{B}}$ the change of coordinates matrix from \mathcal{B} to \mathcal{C} .

$$\begin{aligned}\mathcal{B} &= \{5x^2 - x - 6, 6x^2 + 8x + 9, 3x^2 + 11x + 8\}, \\ \mathcal{C} &= \{x^2 + x + 1, x + 1, 1\}.\end{aligned}$$

Part b. Write the polynomial $(5x^2 - x - 6) + (6x^2 + 8x + 9)$ in terms of the basis \mathcal{C} .

Problem 5 [20 points]

In each of the following cases, clearly mark the statement as **true** or **false**. Please also explain your answers in order to receive credit for this problem.

1. Any square matrix A such that $N(A) = \{\vec{0}\}$ always has an inverse.
2. Let A, B, C be invertible matrices. Then, the product ABC is always invertible.
3. Consider an elementary matrix E that adds two times row 1 to row 2. Then the (2,1) entry of E^{100} is 2^{100} .
4. If a 4x3 matrix has 3 pivots, then $A\vec{x} = \vec{b}$ always has at least a solution.
5. Let V be the set of 2 by 2 matrices. Then, the set of symmetric 2 by 2 matrices is a subspace of V .

6. Consider the set of vectors \vec{x} such that $A\vec{y} = \vec{x}$ always has a solution. Is that a subspace?
7. Let $A = LU$ (L lower triangular with ones on the diagonal, U upper triangular). Let $\vec{a}_1, \vec{a}_2, \vec{a}_3$ be the columns of A and $\vec{u}_1, \vec{u}_2, \vec{u}_3$ be the columns of U . Then, if $\vec{u}_3 = 2\vec{u}_2 - \vec{u}_1$, we also have that $\vec{a}_3 = 2\vec{a}_2 - \vec{a}_1$.
8. If A is a change of basis matrix from a basis $\{\vec{v}_1, \vec{v}_2\}$ to a basis $\{\vec{u}_1, \vec{u}_2\}$, and B is a change of basis matrix from the basis $\{\vec{u}_1, \vec{u}_2\}$ to a basis $\{\vec{w}_1, \vec{w}_2\}$, then AB is a change of basis matrix from $\{\vec{v}_1, \vec{v}_2\}$ to $\{\vec{w}_1, \vec{w}_2\}$.

Extra space for work:

Extra space for work:

Extra space for work:

Extra space for work: